



FUNGICIDE

- **PREVENTATIVE AND CURATIVE CONTROL**
- **Ø** RAINFAST IN 1 HOUR
- **V** REDUCES NEED FOR SWITCHING FOR UP TO 2 WEEKS

A contact acting fungicide for the control of Fusarium Patch (Microdochium nivale), Red Thread (Laetisaria fuciformis) and Dollar spot (Sclerotinia homeocarpa). Useful reduction of Anthracnose (Colletotrichum graminicola), Leaf spot (Drechslera poae) and Rust (Puccinia coronata) where present at time of application.

A suspension concentrate containing 255 g/litre (25% w/w) iprodione.

MAPP 13843

5Le



SAFETY PRECAUTIONS

Operator Protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

TAKE OFF IMMEDIATELY heavily contaminated clothing.

WASH SPLASHES OF CONCENTRATE from skin and eyes immediately.

When using do not eat drink or smoke.

WASH HANDS before meals and after work.

In case of accident or if you feel unwell seek medical advice immediately (show the label where possible).

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

To protect aquatic organisms respect an unsprayed buffer zone to surface water bodies in line with LERAP requirements.

DO NOT ALLOW DIRECT SPRAY from horizontal boom sprayers to fall within 5 m of the top of the bank of a static or flowing water body, unless a Local Environment Risk Assessment for Pesticides (LERAP) permits a narrower buffer zone, or within 1 m of the top of a ditch which is dry at the time of application. DO NOT ALLOW DIRECT SPRAY from hand-held sprayers to fall within 1 m of the top of the bank of a static or flowing water body.

Aim spray away from water.

This product qualifies for inclusion within the Local Environment Risk Assessment for Pesticides (LERAP) scheme. Before each spraying operation from a horizontal boom sprayer, either a LERAP must be carried out in accordance with CRD's published guidance or the statutory buffer zone must be maintained. The results of the LERAP must be recorded and kept available for three years.



Storage and Disposal

Keep out of the reach of children.

Keep away from food, drink and animal feeding stuffs.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.
WASH OUT CONTAINER THOROUGHLY and dispose of safely.

DO NOT RE-USE CONTAINER for any purpose.

PROTECT FROM FROST.

SHAKE THE CONTAINER WELL BEFORE USE

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

CHIPCO® GREEN can be applied year round to managed amenity turf (When used on golf fairways patch treatment of up to 5 % of the total area).

GENERAL INFORMATION:

CHIPCO® GREEN is a contact fungicide containing iprodione which has curative and preventative action against a range of turf diseases. Its special liquid

formulation is 'rainfast' within 1 hour and can prevent the build-up of morning dew on grass leaves. 'Switching' to remove this dew is therefore usually unnecessary.

RESTRICTIONS:

If grass is being mown, CHIPCO® GREEN should be applied after cutting and further mowing should not take place until AT LEAST 24 hours after treatment.

APPLICATION RATE:

For control of Red Thread, Fusarium Patch and Dollar spot. Useful reduction of Anthracnose, Rust and Leaf spot where present at time of application.

Application Equipment	Vehicle-mounted Hydraulic Sprayer		Knapsack Sprayer	
Disease Problem	Rate per ha	Water per ha	Rate per 100 m²	Water per 100m m²
Microdochium nivale (Fusarium Patch), Laetisaria fuciformis (Red Thread) Colletotrichum graminicola (Anthracnose)* Puccinia coronata (Rust)* Sclerotinia homeocarpa (Dollar spot) Drechslera poae (Leaf spot)*	20 L	500 L	200 ml	5 L

^{*}useful reduction for when present at time of application against the other listed diseases.

TIMING OF APPLICATION:

Curative: Apply at the first sign of disease and repeat as necessary at intervals of 1 month or greater.

Preventative: Apply at monthly intervals.

CHIPCO® GREEN can be applied year round to golf courses.

MIXING AND SPRAYING:

Add the appropriate quantity of CHIPCO® GREEN to the spray tank, which contains half the required volume of water. Mix thoroughly and make up to the required volume with water. Maintain adequate agitation during application. Continue agitation at all times during spraying and stoppages until the tank is completely empty. Spray immediately after mixing.

Wash out all spray equipment thoroughly with water immediately after use.

Resistance Management:

CHIPCO® GREEN belongs to the dicarboxamide class of chemistry. There has been no record of managed amenity turf pathogens developing resistance to CHIPCO® GREEN.

To avoid the development of resistance, a resistance management strategy should be adopted. This strategy should include using in rotation and/or tank mixture with products that have a different mode of action or different chemical group that are effective against the target disease, as well as non chemical methods of control. Bayer recommends responsible product stewardship to ensure effective long term control of the fungal diseases on this label.

Important Information

FOR USE ONLY AS AN AMENITY/ HORTICULTURAL FUNGICIDE

Situation Managed amenity turf

Maximum individual dose: 20 L/ha
Maximum number of treatments: 4 per year

Other specific Restrictions

i) For use on (a) commercial and residential lawns up to 200 m² and patch treatment up to 5% of estate lawns of up to 5000 m² (i.e. 250 m²) (b) bowling greens, cricket pitches (excluding outfield), lawn tennis courts and professional sports grounds/stadiums (c) golf greens, tees and patch treatment of up to 5% on fairways'

ii) A minimum of four weeks must be observed between treatments.

Read the label before use. Using this product in a manner that is inconsistent with the label may be an offence. Follow the code of practice for using plant protection products.

Safety Information

CHIPCO® GREEN

A suspension concentrate containing 255 g/Litre (25%w/w) iprodione.





Warning

Suspected of causing cancer.

Very toxic to aquatic life with long lasting effects.

Wear protective gloves/protective clothing/eye protection/face protection.

IF exposed or concerned: Get medical advice/attention.

Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste.

Contains 1,2-benzisothiazolin-3-one. May produce an allergic reaction.

To avoid risks to human health and the environment, comply with the instructions for use.

MAPP 1384

Chipco Green - GB - 5 L - GB80953801F - ARTICLE 80897464 - A3b

For 24 hour emergency information contact Bayer CropScience Limited Telephone: 0800 220876

Batch number : See on the pack

Bayer Environmental Science
Bayer CropScience Limited
230 Cambridge Science Park
Milton Road
Cambridge

CB4 0WB

Tel: 00800 1214 9451

ntal Science. www.environmentalscience.bayer.co.uk for MSDS and larger label

CHIPCO® is a registered trademark of Bayer Environmental Science.